



GB00/3573

10/088945



The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

4




I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 21 September 2000

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)



22SEP99 E478508-1 D02093  
P01/7700 0.00 - 9922346.3


The Patent Office

Cardiff Road  
Newport  
Gwent NP23 5JH

21 SEP 1999

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



1. Your reference

PPD 50449/GB/P

2. Patent application number

(The Patent Office will fill in this part)

9922346.3

3. Full name, address and postcode of the or of each applicant (underline all surnames)

ZENECA Limited  
15 Stanhope Gate  
London W1Y 6LN  
UNITED KINGDOM

Patents ADP number (if you know it)

6254007002

If the applicant is a corporate body, give the country/state of its incorporation

UNITED KINGDOM

4. Title of the invention

IMPROVEMENTS IN OR RELATING TO  
ORGANIC COMPOUNDS

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Frank Mackie HUSKISSON  
Intellectual Property Department  
ZENECA Agrochemicals  
Jealott's Hill Research Station  
P O Box 3538  
Bracknell Berkshire RG42 6YA  
UNITED KINGDOM

Patents ADP number (if you know it)

03989060002

6634547001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 38

Claim(s) 5

Abstract —

Drawing(s) 3 + 3

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77) X

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

ZENECA LIMITED

Signature

*[Signature]*

Date 21/09/99

Authorised Signatory

12. Name and daytime telephone number of person to contact in the United Kingdom

VIJAYA KUMARI MALLIPEDDI  
DAWN LISA ROYAL

01344 414365  
01344 414079

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

IMPROVEMENTS IN OR RELATING TO ORGANIC COMPOUNDS

The present invention relates *inter alia*, to polynucleotide sequences and their use in methods of providing herbicide resistance in plants. In particular the polynucleotides and methods may be used to confer resistance to herbicides comprising a chemical selected from the group consisting of: fomesafen; acifluorfen; chlorimuron ethyl and acetochlor.

Plants which are substantially "tolerant" to a herbicide when they are subjected to it provide a dose/response curve which is shifted to the right when compared with that provided by similarly subjected non tolerant like plants. Such dose/response curves have "dose" plotted on the x-axis and "percentage kill", "herbicidal effect" etc. plotted on the y-axis. Tolerant plants will typically require at least twice as much herbicide as non tolerant like plants in order to produce a given herbicidal effect. Plants which are substantially "resistant" to the herbicide exhibit few, if any, necrotic, lytic, chlorotic or other lesions when subjected to the herbicide at concentrations and rates which are typically employed by the agrochemical community to kill weeds in the field. Hereinafter the words (i) "tolerant" and (ii) "resistant" when used individually mean "tolerant and/or resistant".

Herbicide resistant plants are already available within the art for example, ROUNDUP READY™ Soya which is resistant to herbicides having as a site of action the enzyme 5-enolpyruvylshikimate-3-phosphate synthase, such as those agrochemicals containing glyphosate. One of the advantages of these plants is that the farmer can apply the herbicides to fields containing the resistant crop plants and weeds using "over-the-top application", to kill the weeds.

Other examples of herbicide resistant plants and products and methods for their production are shown in International Patent Application Publication Number WO 93/01294 and WO99/14337. Here the resistance is achieved by inserting into the plant a polynucleotide which provides for the production of a glutathione-S-transferase (GST) enzyme which is involved with the detoxification of the herbicide. Glutathione-S-transferase enzymes have been shown to exist in various organisms such as bacteria, fungi, yeast, plants, mammals and fish and may exist as homo or heterodimers with subunits typically between 24 and 30 KDa. It has been shown that herbicide detoxification is achieved by the conjugation of the herbicide with the free thiol glutathione (GSH), a tripeptide (gamma-

glutamyl-cysteinyl-glycine) within the plant (Cole D.J. 1994 Pesticide Science. 42 pp209-222). Such conjugation is catalysed by GST. Detoxification of herbicides has also been shown to occur following the conjugation of the herbicide with homoglutathione, which is the predominant thiol in some leguminous species. Homoglutathione (hGSH) is also a

5 tripeptide (gamma-glutamyl-cysteinyl-Beta-alanine) but differs from GSH by the addition of Beta-alanine instead of a glycine to the gamma-glutamyl-cysteinyl part.

Thus, the present invention seeks to provide *inter alia*, novel polynucleotides which encode proteins which can be used in methods of providing herbicide resistant plants.

10 According to the present invention there is provided a protein comprising the amino acid sequence depicted as SEQ ID No. 1 or a protein variant having at least about 70% identity therewith wherein the said protein or variant is capable of catalysing the addition of Beta-alanine onto gamma glutamylcysteine. In particular the protein variant may be achieved through conservative substitutions within the amino acid sequence of the active  
15 protein which substitutions do not significantly adversely affect the activity of the variant protein. For example substitutions may be made between the following amino acid groups *viz.*

- (a) Alanine, Serine, Glycine and Threonine
- (b) Glutamic acid and Aspartic acid
- 20 (c) Arginine and Lysine
- (d) Isoleucine, Leucine, Valine and Methionine
- (e) Phenylalanine, Tyrosine and Tryptophan

It is particularly preferred that the protein variant has a  $K_m$  for Beta-alanine which is less than the said protein variants  $K_m$  for glycine when calculated using the same method.

25 Preferably, the  $K_m$  for Beta-alanine is less than or equal to about 10mM, more preferably 5mM, more preferably 4mM, more preferably 3mM, more preferably 2mM, more preferably 1mM and more preferably 0.8mM. The comparison of the  $K_m$  with respect to Beta-alanine and glycine of the protein variants according to the present invention is calculated using the same method.

30 It is further preferred that the said protein variant comprises a region having the amino acid sequence depicted as SEQ ID No. 2 (KKIQQELAKP) and/or SEQ ID No. 3

(CFAGLWSL) and/or SEQ ID No. 4 (VMKPQREGGGNNIYG) and/or SEQ ID No. 5 (AA YILMQRIFP). The present invention still further provides a polynucleotide encoding the protein or protein variant referred to above and it is particularly preferred that the said polynucleotide comprises the sequence depicted as SEQ ID No. 6. Proteins and protein

5 variants falling within this group may be referred to as homoglutathione synthetases.

In a further aspect of the present invention there is provided a Glutathione-S-transferase comprising the amino acid sequence depicted as SEQ ID No. 7 or 8 or 9 or 10 and a polynucleotide comprising a region which encodes at least one of the said amino acid sequences, preferably the sequence depicted as SEQ ID No. 11 or 12 or 13 or 14. The  
10 present invention also provides a polynucleotide sequence which is the complement of one which will hybridise to the polynucleotide of the preceding sentence under stringent conditions and which polynucleotide sequence still encodes a Glutathione-S-transferase. An example of such stringent conditions is hybridisation conducted at 65°C in a solution containing 6xSSC, 0.01% SDS and 0.25% skimmed milk powder, followed by washing at 65  
15 °C in a solution containing 0.2xSSC and 0.1% SDS.

In a further aspect of the present invention there is provided a polynucleotide sequence which is the complement of one which binds to SEQ ID No. 11 or 12 or 13 or 14 at a temperature of between 60°C and 65°C in 0.3 strength citrate buffered saline containing 0.1% SDS followed by rinsing at the same temperature with 0.3 strength citrate buffered  
20 saline containing 0.1% SDS wherein said polynucleotide sequence still encodes a functional Glutathione-S-transferase enzyme with the *proviso* that the said polynucleotide sequence or the amino acid sequence which it encodes, is not a sequence selected from the group of sequences listed under accession number; P32110, U20809, Q03663, P32111, P46421, AJ000923, AF004358, AC000348 or AF051238. In particular the polynucleotide sequence  
25 of the preceding sentence is not a sequence selected from the group depicted as SEQ ID Nos. 25 to 34.

The present invention further provides a DNA construct comprising in sequence a plant operable promoter operably linked to a polynucleotide or polynucleotide sequence encoding a GST according to the present invention operably linked to a terminator region.

The present invention still further provides a DNA construct comprising in sequence a plant operable promoter operably linked to a polynucleotide or polynucleotide sequence encoding a protein or protein variant referred to above operably linked to a terminator region.

The present invention still further provides a DNA construct comprising a first region  
5 comprising in sequence a plant operable promoter operably linked to a polynucleotide encoding a protein or protein variant referred to above and a terminator region and a second region comprising in sequence a plant operable promoter operably linked to a polynucleotide or polynucleotide sequence encoding a GST according to the present invention and a terminator region. It is particularly preferred that the said first region contains a  
10 polynucleotide encoding the amino acid sequence depicted as SEQ ID No. 1 and the said second region contains a polynucleotide encoding the amino acid sequence depicted as SEQ ID No. 10. It is also preferred that the DNA construct referred to above further comprises a third region which provides for the production of a protein which acts as a selectable marker such as those that provide for antibiotic resistance e.g. kanamycin resistance or those  
15 providing for herbicide resistance e.g. glufosinate resistance.

It is preferred that the plant operable promoter within the DNA construct according to the present invention is constitutive, tissue specific, inducible or developmentally regulated. Such promoters, which are *per se* not germane to the invention, are well known to the skilled person and include, for example such as the constitutive CaMV35S, FMV35S, NOS, OCS,  
20 Patatin and E9 (derived from the small subunit of RUBISCO) or inducible such as the alcA/alcR gene switch described in published International Patent Application No. WO93/21334; the GST promoter switch described in published International Patent Application Nos. WO90/08826 and WO93/01294 and the RMS switch system described in published International Patent Application No. WO90/08830. Also the promoter may be  
25 developmentally regulated and/or tissue specific promoters for example the oleosin, ribulose biphosphate carboxylase-oxygenase small sub-unit promoters. Terminators which can be used in the present invention include; Nos and the terminator of a gene of alpha-tubulin (EP-A 652,286).

The DNA construct of the present invention may further comprise (a) transcriptional  
30 enhancing elements; and/or (b) regions encoding non translated translational enhancing sequences, preferably Omega and Omega prime; and/or (c) regions encoding non translated



sequences such as intron sequences; and/or (d) regions encoding target sequences which are capable of directing transcription products to either intracellular organelles, intracellular compartments, cell membranes or to the outside of the cell.

---

In a further aspect of the present invention there is provided a method of providing  
5 plants which are resistant and/or tolerant to an agrochemical comprising (a) inserting into the genome of plant a polynucleotide or a polynucleotide sequence and/or a DNA construct which contains a GST according to the present invention; and (b) regenerating morphologically normal fertile plants or plant parts therefrom; and (c) applying to said plants or plant parts an amount of said agrochemical which is phytotoxic to control like plants and  
10 selecting those plants or plant parts which are resistant to said agrochemical.

In a further aspect of the present invention there is provided a method of providing plants which are resistant and/or tolerant to an agrochemical comprising (a) inserting into the genome of a plant which plant provides for the production of a functional Glutathione-S-transferase, a polynucleotide or a DNA construct comprising a protein or protein variant  
15 according to the present invention; and (b) regenerating morphologically normal fertile plants or plant parts therefrom; and (c) applying to said plants or plant parts an amount of said agrochemical which is phytotoxic to control like plants and selecting those plants or plant parts which are resistant to said agrochemical.

In a further aspect of the present invention there is provided a herbicide resistant  
20 plants obtainable by the method of the preceding paragraph.

In a further aspect of the present invention there is provided a method of providing plants which are resistant and/or tolerant to an agrochemical comprising (a) inserting into the genome of a plant a DNA construct comprising a first region comprising in sequence a plant operable promoter operably linked to a polynucleotide encoding a protein or protein variant  
25 referred to above and a terminator region and a second region comprising in sequence a plant operable promoter operably linked to a polynucleotide or polynucleotide sequence encoding a GST according to the present invention and a terminator region. It is particularly preferred that the said first region comprises a polynucleotide encoding the amino acid sequence depicted as SEQ ID No. 1 and the said second region comprises a polynucleotide encoding  
30 the amino acid sequence depicted as SEQ ID No. 10; and (b) regenerating morphologically normal fertile plants or plant parts therefrom; and (c) applying to said plants or plant parts an

amount of said agrochemical which is phytotoxic to control like plants and selecting those plants or plant parts which are resistant to said agrochemical.

In a further aspect of the present invention there is provided a herbicide resistant plants obtained by the method of the preceding paragraph.

---

5 It is particularly preferred that the agrochemical used in the methods described above comprises acifluorfen and/or chlorimuron-ethyl and/or fomesafen and/or acetochlor and/or metolachlor.

The polynucleotides and polynucleotide sequences and DNA constructs described above may be inserted into the genome of the selected plants using standard transformation techniques including particle mediated biolistic transformation, *Agrobacterium*-mediated  
10 transformation, protoplast transformation (optionally in the presence of polyethylene glycols); sonication of plant tissues, cells or protoplasts in a medium comprising the polynucleotide or vector; micro-insertion of the polynucleotide or vector into totipotent plant material (optionally employing the known silicon carbide "whiskers" technique),  
15 electroporation and the like.

In a further aspect of the present invention there is provided plants or plant parts obtained according to the methods described above. It is particularly preferred that the plants or plant parts are selected from the group consisting of: melons, mangoes, soybean, cotton, tobacco, sugarbeet, oilseed rape, canola, flax, sunflower, potato, tomato, alfalfa, lettuce,  
20 maize, wheat, sorghum, rye, bananas, barley, oat, turf grass, forage grass, sugar cane, pea, field bean, rice, pine, poplar, apple, peaches, grape, strawberries, carrot, lettuce, cabbage, onion, citrus or nut plants.

In a further aspect of the present invention there is provided the use of a polynucleotide or a polynucleotide sequence according or a DNA construct according to the  
25 present invention in a method of producing plants which are resistant and/or tolerant to a herbicide comprising acifluorfen and/or chlorimuron-ethyl and/or fomesafen and/or acetochlor and/or metolachlor.

In a further aspect of the present invention there is provided a method of selectively controlling weeds in a field said field comprising crop plants and weeds said method  
30 comprising applying to said field an agriculturally acceptable formulation of an agrochemical comprising acifluorfen and/or chlorimuron-ethyl and/or fomesafen and/or acetochlor and/or

metolachlor or a functionally related analogue thereof, characterised in that the said crop plants are the plants obtained according to the methods described above.

In a further aspect of the present invention there is provided the use of an agrochemical comprising acifluorfen and/or chlorimuron-ethyl and/or fomesafen and/or

5 acetochlor and/or metolachlor or a functional analogue thereof to selectively control weeds in a field which field comprises crop plants and weeds comprising applying to said field an agriculturally acceptable formulation of said agrochemical in an amount which is sufficient to be phytotoxic to said weeds but not said crop plants characterised in that said crop plants are the plants obtained according to the methods described above.

10 In a further aspect of the present invention there is provided a protein comprising the sequence depicted as SEQ ID No. 7 or a protein variant having a Smith-Waterman score greater than 1094 in the SWISSPROT database calculated using the FASTA3 algorithm wherein the said protein variant still encodes a glutathione-S-transferase.

15 In a further aspect of the present invention there is provided a protein comprising the sequence depicted as SEQ ID No. 8 or a protein variant having a Smith-Waterman score greater than 619 in the SWISSPROT database calculated using the FASTA3 algorithm wherein the said protein variant still encodes a Glutathione-S-transferase.

20 In a further aspect of the present invention there is provided a protein comprising the sequence depicted as SEQ ID No. 9 or a protein having a Smith-Waterman score greater than 671 in the SWISSPROT database calculated using the FASTA3 algorithm wherein the said protein variant still encodes a Glutathione-S-transferase.

25 In a further aspect of the present invention there is provided a protein comprising the sequence depicted as SEQ ID No. 10 or a protein having a Smith-Waterman score greater than 766 in the SWISSPROT database calculated using the FASTA3 algorithm wherein the said protein variant still encodes a Glutathione-S-transferase.

30 The Fasta algorithm referred to above is well known to the skilled artisan and uses the method of Pearson and Lipman (Proc. Natl. Acad. Sci. USA 85; 2444-2448 (1988)) to search for similarities between one sequence and any group of sequences of the same type as the query sequence. Fasta also determines the best segment of similarity between the query sequence and the sequences in the database, using a variation of the Smith-Waterman algorithm. This "local alignment" procedure is described in Chao, Pearson, and Miller

(CABIOS 8; 481-487 (1992)). The score for this alignment is reported as the *opt* and Smith-Waterman score. The database used in the above calculations is the SWISSPROT database. This database, which is well known and frequently used by the person skilled in the art is commercially available from sources such as Geneva Bioinformatics (GeneBio™) S.A 25  
5 avenue de Champel CH - 1206 Geneva Switzerland.

The present invention will now be described by way of the following non limiting examples with reference to the figures and sequence listing of which:

Figure 1. Vector pMJB2.

10 Figure 2. Vector pMOG800.

Figure 3. Vector pMOG1051.

#### Sequence Listing

SEQ ID No. 1. Homoglutathione synthetase from *Glycine max*.

15 SEQ ID Nos. 2 to 5. Homoglutathione synthetase protein fragments.

SEQ ID No. 6. Polynucleotide sequence encoding Homoglutathione synthetase from *Glycine max*.

SEQ ID Nos. 7 to 10. Glutathione-S-transferases 2.6, 3.1, 3.2 and 3.3 respectively.

SEQ ID No. 11 to 14. Polynucleotides encoding the GSTs 2.6, 3.1, 3.2 and 3.3 respectively.

20 SEQ ID Nos. 15 to 24. Primers.

SEQ ID No. 25 Soybean sequence P32110 derived nucleic acid sequence.

SEQ ID No. 26 Mungbean sequence U20809.

SEQ ID No. 27 Tobacco sequence Q03663.

SEQ ID No. 28 Potato sequence P32111 derived nucleic acid sequence.

25 SEQ ID No. 29 Arabidopsis sequence P46421.

SEQ ID No. 30 Arabidopsis sequence P46421 (genomic).

SEQ ID No. 31 Papaya sequence AJ000923.

SEQ ID No. 32 Spruce sequence AF051214.

SEQ ID No. 33 Wheat sequence AF004358.

30 SEQ ID No. 34 Spruce sequence AF051238.

Example

Production of plants with increased herbicide tolerance through increased *in planta* expression of (a) homoglutathione (b) homoglutathione and glutathione *S*-transferase (GST).

---

5    1.1 Isolation and preparation of polynucleotide sequence encoding hGSH synthetase.

Degenerate oligonucleotide primer MS-3 and OG9 are used in conjunction with PCR to amplify polynucleotide sequences encoding partial length Homoglutathione (hGSH) synthetase from first strand cDNA, synthesised using total RNA extracted from soybean cell cultures (cv. Mandarin) (Skipsey *et al.* 1997)

10

SEQ ID No. 15

MS-3            5' GCG AAG CCH CAR MGA GAR GGH GGA GG.3'

SEQ ID No. 16

15    OG-9            5' CGC ACT GAG AGA GGA TCC TCG AG 3'

The resulting PCR products are cloned into the vector pGEM-T (Promega™), which is termed MS3-1. The cloned insert is then excised from MS3-1 using *Eco* R1, labelled with digoxigenin using standard protocols (Boehringer Mannheim™), and used to screen a  
20    soybean cell culture cDNA library (cv. Mandarin), constructed in the uni-ZAP XR vector (Stratagene™).

1.2 Isolation and preparation of polynucleotide sequences encoding glutathione *S*-transferase

Degenerate oligonucleotide primers CJACON2 and CJACON3 are used independently in  
25    conjunction with oligonucleotide primer OG9 to amplify DNA probes from first strand cDNA suitable for cDNA library screening.

SEQ ID No. 17.

CJACON2    5' TTC TGG GYK RAS TWC ITY GAC RAI AAG 3'

30

SEQ ID No. 18.

CJACON3 5' GAG TCY MWK GTS ATT GTT GAA TAC ATT GAT GAG 3'

---

5 The PCR products obtained are cloned into the vector pCR 2.1 using the original TA cloning kit (Invitrogen™). Distinct cDNAs are identified by automated DNA sequencing (ABI 377) and labelled with digoxigenin in order to facilitate the screening of cDNA libraries for cDNAs encoding full length enzymes.

### 1.3 Preparation of constructs for use in heterologous gene expression in *E. coli*

10 Full length cDNAs encoding homogluthathione synthetase and glutathione *S*-transferase are independently expressed in *E. coli* using the pET expression system (Novagen™). *Nco* 1 and *Xho* 1 restriction enzyme sites are introduced by at the 5 prime and 3 prime ends of the homogluthathione synthetase cDNA respectively using primers MS4-*Nco* and MS-4-HIS.

15 SEQ ID No. 19

MS4-*Nco* 5' CCT CTC AAA CCC ATG GCT CAA CC 3'

SEQ ID No. 20

MS-4-His 5' GCG CTC GAG AGT TAG GTA TAC AGT ATC TAC C 3'

20

The PCR fragment is then digested with *Nco*1 and *Xho* 1 and ligated into similarly digested pET-24d. This vector is termed pET-MS4-His.

With respect to the cDNAs encoding glutathione *S*-transferase, either *Nde* I or *Nco* I sites are introduced as appropriate at the 5' end, and *Bam* H1 at the 3' end using PCR as described  
25 previously. The cDNA is then introduced into pET-24a or pET-24d as appropriate.

### 1.4 Expression and characterisation of recombinant enzymes in *E. coli*

#### Glutathione *S*-transferase

30 The vectors described in section 1.3, harbouring cDNAs encoding GSTs, are transformed into *E. coli* BL21 (DE3) using standard bacterial transformation procedures. Expression and

purification of the recombinant GST is performed according to the methods described by Cummins *et al.*, 1997.

---

#### Homogluthathione synthetase

- 5 Purification of homogluthathione synthetase is performed using the following method. *E. coli* BL21 (DE3) harbouring the pET-MS4-His plasmid (section 1.3) are grown at 30 °C until  $OD_{600}=0.5$ , after which isopropyl- $\beta$ -D-thiogalactoside (IPTG) is added to a final concentration of 0.1 mM. Following a 3 hour incubation the bacteria are collected by centrifugation, re-suspended in buffer A (20mM Tris, 0.5 M NaCl, 5 mM imidazole pH 8.0)
- 10 and lysed. Cell debris is removed by centrifugation (10,000 g, 10 min) and the supernatant applied to a 5 ml iminodiacetic acid column (Sigma™), previously charged with  $NiSO_4$  and equilibrated in buffer A. The column is then washed buffer A containing 20 mM imidazole, followed by buffer A containing 300 mM imidazole to remove affinity bound protein. The protein in this fraction is then concentrated using a Centriplus 30 (Amicon™) spin column
- 15 and re-suspended in buffer A prior to application onto a 1 ml HiTrap chelating column (Pharmacia™), pre-charged with  $NiSO_4$  and equilibrated as described previously. Affinity bound protein is recovered using an increasing concentration of 20-200 mM imidazole and the presence of His-tagged recombinant homogluthathione synthetase detected using His-tagged antibodies according to standard procedures. Fractions containing recombinant
- 20 homogluthathione synthetase are pooled, concentrated using Centricon 30 spin columns (Amicon™) and resuspended in 20 mM Tris-HCl pH 8.0, 1mM DTT.

#### 1.5 Assay of enzyme activity of recombinant GST and homogluthathione synthetase.

GST assays are performed as described by Andrews *et al.*, 1997.

25

- Homogluthathione synthetase is assayed for activity in 250 mM Tris-HCl pH 8.0, 50 mM KCl, 20 mM  $MgCl_2$ , 5 mM DTT, 10 mM ATP, 1 mM  $\gamma$ -glutamylcysteine and 50 mM glycine or 10 mM  $\beta$ -alanine in a total 100  $\mu$ l. Experimental controls consist of the exclusion of enzyme or glycine/ $\beta$ -alanine. Assays are performed at 30°C for 60 min, with 20  $\mu$ l
- 30 aliquots removed at regular time intervals. Monobromobimane derivatisation is then

performed on the aliquot to determine the presence of either glutathione or homoglutathione according to methods described by Cummins *et al.*, 1997.

---

1.6 Production of plant transformation vector harbouring cDNA encoding homoglutathione synthetase.

Oligonucleotide primers hGSH-Nco1 and hGSH-Kpn1 are used in conjunction with PCR to introduce *Nco*1 and *Kpn*1 sites at the 5 prime and 3 prime end of the cDNA respectively.

SEQ ID No. 21

10 Hgsh-Nco1 5' CCT CTC AAA CCC ATG GCT CAA CC 3'

SEQ ID No. 22

hGSH-Kpn 1 5' CGC GGT ACC TCC ATA CAA AGA AAA TCA 3'

15 The cDNA is then inserted into the vector pMJB2 (Figure 1) as a *Nco*1/*Kpn*1 fragment. The expression cassette, containing the CaMV35S double enhancer : Glucanase II leader, hGSH synthetase cDNA and *nos* terminator is the excised from pMJB2 using *Hind* III/ *Eco*R1 and ligated into the similarly digested binary vector pMOG800 (Figure 2).

20 1.7 Production of plant transformation vector harbouring tandem cDNAs encoding homoglutathione synthetase and GST.

Oligonucleotide primers 3.6-Bgl II and 3.6 *Nco* 1 are used in conjunction with PCR to introduce *Nco*1 and *Bgl* II sites at the 5 prime and 3 prime end of the GST 3.6 cDNA respectively.

25

SEQ. ID No. 23

3.6-Bgl II 5' GAG ATC TGC AAC AAA CAT AGC CTC 3'

SEQ ID No. 24

30 3.6 *Nco* 1 5' TAC ACC ATG GCT GAA AGG GAC TTG 3'



The GST cDNA is then inserted into the vector pMOG1051 (Figure 3) as a *Nco*I/*Bgl* II fragment. The expression cassette, containing the RolD/Fd promoter, GST cDNA and potato PI-II terminator is excised from pMOG1051 using *Bam* HI and ligated into the unique *Bam* HI site in the binary vector containing the hGSH expression cassette (section 1.6).

5 Orientation of the insert is determined by PCR.

### 1.8 Plant Transformation and Regeneration.

Constructs from section 1.6 and 1.7 are transformed into *Agrobacterium tumefaciens* strain  
10 LBA 4404 using the freeze thaw method of transformation provided by Holsters *et al.*, 1978. Tobacco transformation and whole plant regeneration is performed using var. Samusun according to protocols detailed by Bevan, 1984. Transformation events are selected on MS-media containing kanamycin.

### 15 1.9 Analysis of Transgenic Plants

#### PCR analysis of transformants

Leaf samples were taken from transformed lines and DNA extracted according to the methods described by Edwards *et al.*, 1991. Oligonucleotide primers are designed to specific regions within the transgene to enable its detection in the plant material tested.

20

#### RNA analysis

The presence of mRNA encoding the transgene is detected within the plant using Northern Blot hybridisation. Total RNA is extracted from leaf tissue using Tri-reagent and protocols provided by the manufacturer (Sigma™). Blots are prepared using existing procedures  
25 (Sambrook, 1989) and probed using radio-labelled cDNA encoding the desired mRNA.

#### Protein Analysis

The presence of recombinant protein in the transgenic plant is determined using Western blotting procedures with antibodies raised to the relevant enzyme using standard protocols.

30

### Herbicide tolerance tests

Following tissue culture, transgenic plants are transferred to 5 inch pots containing John Innes potting compost no. 3. The plants are allowed to acclimatise (approx. 2 weeks) and various herbicides applied at various concentrations to the aerial tissue using a track sprayer.

- 5 Visual assessment of phytotoxicity/plant necrosis is performed 6 and 13 days post application. Plants show resistance to herbicides applied at concentrations which are phytotoxic to control like plants.

### 1.10 Production of homozygous plant lines

- 10 Single copy transgenic plant lines are identified by Southern blot analysis according to methods described by Sambrook, 1989 using appropriate radiolabelled probes. Segregation analysis is performed on plants containing single insertion events by germination on MS media containing kanamycin. Further confirmation of homozygous lines is performed by back crossing transgenic lines with wild-type tobacco and analysis genetic segregation  
15 following selection on kanamycin.

### References

- Andrews C.J., Skipsey M., Townson J.K., Morris C., Jepson I. and Edwards R (1997) Glutathione transferase activities toward herbicides used selectively in soybean. *Pesticide*  
20 *Science* 51 213-222.
- Bevan M.M (1984) Binary *Agrobacterium* vectors used for plant transformation. *Nucleic Acids Research* 12 8711-8721.
- Cummins I., Moss S., Cole D.J and Edwards R (1997a) Glutathione Transferases in  
25 Herbicide-resistant and Herbicide-Susceptible Black-grass (*Alopecurus myosuroides*). *Pesticide Science* 51 244-250.
- Edwards K., Johnstone C and Thompson C (1991) A simple and rapid method for the preparation of plant genomic DNA for PCR analysis. *Nucleic Acids Research* 19 1349.

Holsters M., de Waele D., Depicker A., Messens E., van Montagu M and Schell J (1978) Transfection and transformation of *Agrobacterium tumefaciens*. *Molecular and General Genetics* 163 181-187.

---

- 5 Skipsey M., Andrews CJ, Townson J.K., Jepson I and Edwards R (1997) Substrate and thiol specificity of a stress-inducible glutathione transferase from soybean. *FEBS Letters* 409 370-374.

## SEQUENCE LISTING

&lt;110&gt; ZENECA LIMITED

---

5 <120> IMPROVEMENTS IN OR RELATING TO ORGANIC COMPOUNDS

&lt;130&gt; ZENECA CASE PPD50449

&lt;140&gt;

10 &lt;141&gt;

&lt;160&gt; 34

&lt;170&gt; PatentIn Ver. 2.0

15

&lt;210&gt; 1

&lt;211&gt; 499

&lt;212&gt; PRT

&lt;213&gt; Glycine max

20

&lt;400&gt; 1

Met Ser Gln Pro Leu Thr Thr Asn Ser Val Leu Val Glu Glu Ala Ala  
1 5 10 15

25 Ala Asp Gly Asp Ser Ser Ala Ala Ala Pro Pro Leu Phe Asp Tyr His  
20 25 30

Arg Ile Asp Gln Lys Leu Leu Gln Asn Ile Val Tyr Asp Ala Leu Val  
35 40 45

30

Trp Ser Thr Leu Asn Cys Leu Leu Val Gly Asp Lys Ser Val Gln Arg  
50 55 60

35 Ser Gly Arg Val Pro Gly Val Gly Leu Val His Leu Pro Leu Ser Leu  
65 70 75 80

Leu Pro Gly Pro Phe Pro Glu Ser His Trp Lys Gln Gly Cys Glu Leu  
85 90 95

40 Ala Pro Ile Phe Asn Glu Leu Val Asp Arg Val Ser Leu Asp Gly Lys  
100 105 110

Phe Leu Gln Glu Ser Leu Ser Arg Thr Lys Asn Ala Asp Glu Phe Thr

	115	120	125
	Ser Arg Leu Leu Asp Ile His Ser Lys Met Leu Gln Ile Asn Lys Lys		
	130	135	140
5	<hr/>		
	Glu Asp Ile Arg Met Gly Ile Val Arg Ser Asp Tyr Met Ile Asp Glu		
	145	150	155 160
10	Lys Thr Lys Ser Leu Leu Gln Ile Glu Met Asn Thr Ile Ser Thr Ser		
	165	170	175
	Phe Ala Leu Ile Gly Cys Leu Met Thr Gly Leu His Lys Ser Leu Leu		
	180	185	190
15	Ser Gln Tyr Gly Lys Phe Leu Gly Leu Asn Ser Asn Arg Val Pro Ala		
	195	200	205
	Asn Asn Ala Val Asp Gln Ser Ala Glu Ala Leu Ala Lys Ala Trp Ser		
	210	215	220
20	Glu Tyr Asn Asn Pro Arg Ala Ala Ile Leu Val Val Val Gln Val Glu		
	225	230	235 240
	Glu Arg Asn Met Tyr Glu Gln His Tyr Ile Ser Ala Leu Leu Arg Glu		
25	245	250	255
	Lys His His Ile Arg Ser Ile Arg Lys Thr Leu Thr Glu Ile Asp Gln		
	260	265	270
30	Glu Gly Lys Ile Leu Pro Asp Gly Thr Leu Ser Val Asp Gly Gln Ala		
	275	280	285
	Ile Ser Val Val Tyr Phe Arg Ala Gly Tyr Thr Pro Lys Asp Tyr Pro		
	290	295	300
35	Ser Glu Ser Glu Trp Arg Ala Arg Leu Leu Met Glu Gln Ser Ser Ala		
	305	310	315 320
	Ile Lys Cys Pro Thr Ile Ser Tyr His Leu Val Gly Thr Lys Lys Ile		
40	325	330	335
	Gln Gln Glu Leu Ala Lys Pro Gly Val Leu Glu Arg Phe Val Glu Asn		
	340	345	350

Lys Asp His Ile Ala Lys Leu Arg Ala Cys Phe Ala Gly Leu Trp Ser  
 355 360 365

---

5 Leu Glu Asp Ser Asp Ile Val Lys Lys Ala Ile Glu Asn Pro Glu Leu  
 370 375 380

Phe Val Met Lys Pro Gln Arg Glu Gly Gly Gly Asn Asn Ile Tyr Gly  
 385 390 395 400

10

Asp Glu Leu Arg Glu Thr Leu Leu Lys Leu Gln Glu Ala Gly Ser Gln  
 405 410 415

15

Glu Asp Ala Ala Tyr Ile Leu Met Gln Arg Ile Phe Pro Ala Thr Ser  
 420 425 430

Pro Ala Ile Leu Val Arg Asp Gly Asn Trp Asp Thr Gly His Val Ile  
 435 440 445

20

Ser Glu Ala Gly Ile Phe Gly Thr Tyr Leu Arg Asn Lys Asp Lys Ile  
 450 455 460

Ile Ile Asn Asn Glu Ser Gly Tyr Met Val Arg Thr Lys Ile Ser Ser  
 465 470 475 480

25

Ser Tyr Glu Gly Gly Val Leu Pro Gly Phe Gly Val Val Asp Thr Val  
 485 490 495

Tyr Leu Thr

30

<210> 2

<211> 10

35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Protein  
 Fragment

40

<400> 2

Lys Lys Ile Gln Gln Glu Leu Ala Lys Pro

1 5 10

<210> 3

5 <211> 8

<212> PRT

<213> Artificial Sequence

<220>

10 <223> Description of Artificial Sequence:Protein  
Fragment

<400> 3

Cys Phe Ala Gly Leu Trp Ser Leu

15 1 5

<210> 4

<211> 15

20 <212> PRT

<213> Artificial Sequence

<220>

25 <223> Description of Artificial Sequence:Protein  
Fragment

<400> 4

Val Met Lys Pro Gln Arg Glu Gly Gly Gly Asn Asn Ile Tyr Gly

30 1 5 10 15

<210> 5

<211> 11

<212> PRT

35 <213> Artificial Sequence

<220>

40 <223> Description of Artificial Sequence:Protein  
Fragment

<400> 5

Ala Ala Tyr Ile Leu Met Gln Arg Ile Phe Pro

1 5 10

&lt;210&gt; 6

&lt;211&gt; 1854

5 &lt;212&gt; DNA

&lt;213&gt; Glycine max

&lt;400&gt; 6

10 ggcggcttgg tttgttctac ttctctttac actgggatta gaacaaggcg tttgtgcact 60  
tctaacaaca ccacctttcc cttcccccaa caacctcaat cactctcttt cgctaaacct 120  
ctcaaactca tgtctcaacc tttgaccacc aactctgttc ttgttgaaga ggctgctgct 180  
gatggtgatt cctccgccgc cgcacctccc ctcttcgatt atcatcgtat cgaccaaaaa 240  
ctgctccaaa acatagttaa cgatgctctt gtctggagca ccctcaactg cctccttggt 300  
ggtgacaaat ctgttcagag atcaggaaga gttcctgggtg tgggcctggg acatctccca 360  
15 ctttccttat tacctgggcc atttctgaa agtcattgga agcaagggtg cgaattagct 420  
cctatattta atgaacttgt tgatcgggtg agtttggatg ggaaatttct ccaggaatct 480  
ctctccagaa ctaagaatgc ggatgaattt acctcaagac ttttagatat tcattctaag 540  
atgctacaga ttaacaaaaa agaggacata cgcatgggaa tagttcgttc agattatatg 600  
attgatgaga agactaaatc acttttacia atagagatga aactatttct cacttcattt 660  
20 gctttgattg gttgtcttat gactggactt cataagagct tactttctca atatggaaaa 720  
ttccttggac taaattccaa tagggttcct gccataatg ccgttgatca gtctgcagag 780  
gccttggcta aagcttggag tgagtataac aatcccaggg ctgcaattct ggtcgtggtt 840  
cagggtgaag aaagaaacat gtacgagcag cattatatat ctgcacttct aagagaaaaag 900  
catcatatta gaagcatacg caaacggtg accgaaattg atcaggaagg aaaaattctg 960  
25 ccagatggaa cactttctgt ggatggacaa gcaatttcag ttgtttactt ccgggctggc 1020  
tacacgccaa aggactatcc ttcagaatca gaatggagag ctaggctact gatggaacaa 1080  
tcttctgcta tcaaatgcc tacaatatct tatcatttgg ttggcaccaa aaagattcaa 1140  
caggaacttg caaagcctgg tgttcttgag aggttcgttg aaaacaaaga ccacattgcc 1200  
aaattgcgtg catgctttgc agggttgtgg agtttggag actcagatat tgtaaaaaaa 1260  
30 gcaattgaaa atccagagct atttgtgatg aagcctcaaa gagaaggagg aggaaacaat 1320  
atztatggtg atgagttgag ggaaaccctc cttaaattac aggaagcagg ttctcaagaa 1380  
gatgcagcat acatccttat gcagaggata tttcccgcca cttctccagc aattttggtg 1440  
cgtgatggta attgggatac gggcatgtc atttcagaag ctggaatatt tggacttat 1500  
ttaaggaata aggacaagat tatcattaat aacgaaagtg gctatatggt gcgtacaaaa 1560  
35 atatcatcat cttatgaagg aggagttttg cctggttttg gagtggtaga tactgtatac 1620  
ctaacttgat ggagctaacc cccaagtta tcaaagcaat tcaaaacatt atgtatgggt 1680  
tatatatcac cactcaagtc tcctcactcc tgattttctt tgtatggagg cattgctgtt 1740  
tcttttaatt gttcctatgg gatggtgtct aattattaac tgtactcaac gacctgtttg 1800  
attctaacca ataaagattg atgaactgtt ctaacaaaaa aaaaaaaaaa aaaa 1854

40

&lt;210&gt; 7

&lt;211&gt; 222

&lt;212&gt; PRT



<400> 7

5 1 5 10 15

Phe Lys Ala Arg Ala Gln Ser Phe Val Ala Lys Arg Lys Asn

210

215

220

&lt;210&gt; 8

5 &lt;211&gt; 235

&lt;212&gt; PRT

&lt;213&gt; Glycine max

&lt;400&gt; 8

10 Met Ala Glu Gln Asp Lys Val Ile Leu His Gly Met Trp Ala Ser Pro  
1 5 10 15

Tyr Ala Lys Arg Val Glu Leu Ala Leu Asn Phe Lys Gly Ile Pro Tyr  
20 25 30

15 Glu Tyr Val Glu Glu Asp Leu Arg Asn Lys Ser Asp Leu Leu Leu Lys  
35 40 45

20 Tyr Asn Pro Val His Lys Lys Val Pro Val Leu Val His Asn Gly Lys  
50 55 60

Ala Ile Ala Glu Ser Met Val Ile Leu Glu Tyr Ile Asp Glu Thr Trp  
65 70 75 80

25 Lys Asp Gly Pro Lys Leu Leu Pro Ser Asp Ser Tyr Lys Arg Ala Gln  
85 90 95

Ala Arg Phe Trp Cys His Phe Ile Gln Asp Gln Leu Met Glu Ser Thr  
100 105 110

30 Phe Leu Val Val Lys Thr Asp Gly Glu Ala Gln Gln Lys Ala Ile Asp  
115 120 125

35 His Val Tyr Glu Lys Leu Lys Val Leu Glu Asp Gly Met Lys Thr Tyr  
130 135 140

Leu Gly Glu Gly Asn Ala Ile Ile Ser Gly Val Glu Asn Asn Phe Gly  
145 150 155 160

40 Ile Leu Asp Ile Val Phe Cys Ala Leu Tyr Gly Ala Tyr Lys Ala His  
165 170 175

Glu Glu Val Ile Gly Leu Lys Phe Ile Val Pro Glu Lys Phe Pro Val

		180		185		190	
	Leu Phe Ser Trp Leu Met Ala Ile Ala Glu Val Glu Ala Val Lys Ile						
	195		200		205		

---

5	Ala Thr Pro Pro His Glu Lys Thr Val Gly Ile Leu Gln Leu Phe Arg				
	210		215		220

	Leu Ser Ala Leu Lys Ser Ser Ser Ala Thr Glu				
10	225		230		235

	<210> 9
	<211> 223
15	<212> PRT
	<213> Glycine max

	<400> 9						
	Met Ala Glu Val Lys Leu His Gly Phe Trp Tyr Ser Pro Tyr Thr Leu						
20	1		5		10		15

	Arg Val Val Trp Thr Leu Lys Leu Lys Asp Ile Pro Tyr Gln Asn Ile				
	20		25		30

25	Glu Glu Asp Arg Tyr Asn Lys Ser Leu Gln Leu Leu Glu Tyr Asn Pro				
	35		40		45

	Val Tyr Lys Lys Thr Pro Val Leu Val His Asn Gly Lys Pro Leu Cys				
	50		55		60

30	Glu Ser Met Leu Ile Val Glu Tyr Ile Asp Glu Ile Trp Ala His Asn						
	65		70		75		80

	Ser Leu Leu Pro Ala Asp Pro Tyr Glu Arg Ala Leu Ala Arg Phe Trp					
35		85		90		95

	Val Lys Tyr Ala Asp Asp Asp Met Phe Ser Ala Val Ile Ala Phe Phe				
	100		105		110

40	Leu Ser Asn Asn Asp Glu Glu Arg Glu Lys Ser Ile Glu Lys Ile Trp				
	115		120		125

	Glu His Leu Arg Val Val Glu Asn Gln Cys Phe Gly Asp Gln Lys Lys
--	---

130                      135                      140  
 Phe Phe Gly Gly Asp Ile Ile Asn Ile Met Asp Ile Ala Phe Gly Ser  
 145                      150                      155                      160

---

5                      165                      170                      175  
 Ile Phe Lys Ile Leu Val Val Ala Glu Asp Ile Leu Asp Ala Lys Val

10                      180                      185                      190  
 Leu Glu Asp Glu Lys Phe Pro His Leu His Ser Trp Tyr Asn Asn Phe

15                      195                      200                      205  
 Lys Asp Val Ala Val Ile Lys Glu Asn Leu Pro Asp His Glu Lys Met

20                      210                      215                      220  
 Val Ala Phe Ala Lys Phe Ile Arg Glu Lys Arg Leu Ala Cys Thr

<210> 10  
 <211> 232  
 <212> PRT  
 <213> Glycine max

<400> 10  
 25 Met Ala Glu Arg Asp Leu Arg Leu Leu Gly Ala Trp Phe Ser Pro Phe  
     1                      5                      10                      15

30                      20                      25                      30  
 Ala Leu Arg Val Gln Ile Ala Leu Asn Leu Lys Gly Leu Asp Tyr Glu

35                      35                      40                      45  
 Val Val Glu Glu Thr Leu Asn Pro Lys Ser Glu Leu Leu Lys Ser

40                      50                      55                      60  
 Asn Pro Val His Lys Lys Ile Pro Val Phe Phe His Gly Asp Lys Val

65                      70                      75                      80  
 Ile Cys Glu Ser Ala Ile Ile Val Glu Tyr Ile Asp Glu Val Trp Ser

85                      90                      95  
 Asn Asn Ala Leu Ser Ile Leu Pro Gln Asn Ala Tyr Asp Arg Ala Asn

Ala Arg Phe Trp Val Ser Tyr Ile Asp Asp Lys Trp Leu Thr Ser Leu

		100		105		110										
	Lys	Ser	Val	Leu	Ala	Thr	Glu	Asp	Asp	Glu	Ala	Lys	Lys	Leu	His	Phe
		115					120					125				

---

5

	Glu	Gln	Ala	Glu	Glu	Val	Leu	Glu	Lys	Val	Glu	Glu	Val	Phe	Asn	Lys
	130						135					140				

10

	Cys	Ser	Glu	Gly	Lys	Ala	Tyr	Phe	Gly	Gly	Asp	Thr	Ile	Gly	Phe	Val
	145				150						155					160

15

	Asp	Ile	Gly	Phe	Gly	Ser	Phe	Leu	Ser	Phe	Ile	Arg	Val	Ser	Glu	Asn
					165					170					175	

20

	Met	Asn	Glu	Arg	Lys	Leu	Leu	Asp	Glu	Thr	Lys	Tyr	Pro	Gly	Leu	Thr
					180				185					190		

25

	Leu	Trp	Ala	Glu	Thr	Phe	Ala	Ala	Asp	Pro	Ala	Val	Lys	Gly	Leu	Leu
			195					200					205			

30

	Pro	Glu	Thr	Glu	Lys	Leu	Val	Glu	Phe	Ala	Lys	Ile	Leu	Gln	Leu	Lys
		210				215						220				

35

	Trp	Ala	Ala	Ala	Ala	Ala	Ala	Lys								
	225					230										

40

<210> 11

<211> 885

<212> DNA

<213> Glycine max

<400> 11

	ctgcaatgtc	ttcaagtcag	gaagaggtga	cccttttggg	agttgtggga	agcccatttc	60
35	tacacagggg	tcagattgct	ctcaagttga	agggagttga	atacaaata	ttggaagacg	120
	atttgaacaa	caagagtgat	ttgctcctca	agtataaccc	agtttacaaa	atgattccag	180
	tgcttggtca	caatgagaag	cccatttcag	agtccttgt	gattgttgag	tacattgatg	240
	acacatggaa	aaacaatccc	atcttgcctt	ctgatcccta	ccaaagagcc	ttggctcggt	300
	tctgggctaa	gttcattgat	gacaagtgtg	tggttccagc	atggaaatct	gcttttatga	360
40	ctgatgagaa	agagaaagag	aaggctaaag	aagagttatt	tgaggctctg	agttttcttg	420
	agaatgagtt	gaagggcaag	ttttttggtg	gagaggagtt	tggtttgtg	gatattgctg	480
	ctgtgttaat	acctataatt	caagagatag	cagggttgca	attgttcaca	agtgagaaat	540
	tcccaaagct	ctctaaatgg	agccaagact	ttcacaacca	tccagttgtc	aacgaagtta	600

tgcctcctaa ggatcaactt tttgcctatt tcaaggctcg ggctcaaagc ttcggttgcta 660  
 aaagaaagaa ttaatatagt gagactcaga atttccatcg aggtttcagt attgtatgaa 720  
 atgaaagcta cttgtctatg tttcgttatt gcggttgat tttcattttt caatgaatta 780  
 tgtgatatag gatttctcca tgtcaaaaga tagttcaatt caatcaataa aataaacgaa 840

5 tgagtcgtgt tagagcaaaa aaaaaaaaaa aaaaaaaaaa aaaaa 885

<210> 12

<211> 899

<212> DNA

10 <213> Glycine max

<400> 12

atggcagagc aagacaaggt gatcctacac gggatgtggg ccagccctta tgccaagagg 60  
 gtggaattgg cccttaattt taagggcata ccctatgagt atgttgaaga agacttgaga 120  
 15 aataagagtg atttgcttct aaagtacaac cctgttcaca agaaggttcc tgtacttggt 180  
 cataatggaa aggccattgc tgaatccatg gtgatccttg agtatattga tgaaacatgg 240  
 aaagatggtc ctaaactgct tccaagtgat tcttacaac gagcccaagc tcgattctgg 300  
 tgtcatttca tccaggatca gttaatggag agcactttc tagtagtcaa aactgatgga 360  
 gaagcacaac aaaaggccat tgaccacgtg tatgagaaac tgaaagtgct agaagatgga 420  
 20 atgaagacct atctgggaga aggcaatgct attatctctg gtgttgaaaa caactttgga 480  
 atccttgaca ttgtgttttg tgctttatat ggtgcctaca aggctcatga agaagttatt 540  
 ggctcaagt tcatagtgcc agaaaagttt cctgtgttgt tttcttggtt gatggctatt 600  
 gctgaggttg aagctgtgaa aattgcaact cctccacatg aaaaaacagt gggaattctt 660  
 cagttgttca ggctgtctgc actgaaatct tcttctgcca cagaatgata tatacttcaa 720  
 25 cactttaata gactgtccat cgtttgcttc ttctgagagt ctttaatgta tgtatctttc 780  
 aataacagga tgagtaacac ctgagtatgt aaagcgtgat gatatagaga tatacctcta 840  
 tatatcaaat actcttctat aaacacttct ttctttcctt aaaaaaaaaa aaaaaaaaaa 899

<210> 13

30 <211> 840

<212> DNA

<213> Glycine max

<400> 13

35 atggcagagg tgaagcttca tggattttgg tatagtcctt acactttgag ggtggtatgg 60  
 accttaaagt taaaggatat accatatcaa aacatagaag aagaccgcta caataagagt 120  
 cttcaacttc ttgaatacaa ccagtatcac aagaaaactc cagtgttgtt ccataatgga 180  
 aaacccttat gtgagtccat gcttattgtt gaatacattg atgagatttg ggcacataat 240  
 tcattacttc ctgctgatcc ctacgagaga gctctggcaa ggttttgggt taaatatgct 300  
 40 gatgatgaca tgttttctgc agttattgca ttcttcctta gcaataatga tgaagagcga 360  
 gaaaagagca tagagaagat atgggagcat ctgagggttg ttgagaatca gtgttttggg 420  
 gatcagaaga aatttttttg gggagacatt attaacatta tggacatagc ttttgggtcc 480  
 atattcaaaa ttcttgtggg tgcagaagat attcttgacg cgaaggtcct ggaagatgag 540

```

aaattccctc acttgcatte atgggtataat aatttcaagg atgttgcaagt tattaaagaa 600
aacctcccag accatgagaa aatgggtggct ttgctaagt ttattagaga aaaacgtttg 660
gcatgtacct aagaaagtaa tcttatatga gatcaagtat gaatcacttt gtatctgtct 720
gaatcgtttt gttatgcgtg tttctttagt ttccactcca ttattaggat gtcttgacat 780
5 atctgtgaaa gcaataaaaag tttaatggga tgtactggat taaaaaaaaa aaaaaaaaaa 840

```

<210> 14

<211> 918

<212> DNA

10 <213> Glycine max

<400> 14

```

cataaaactc cacatttctt gctgagtaac ctaacaaaac aaacacaata ttgctccgtg 60
tttgacctgt tatagtaaac agtgatggct gaaagggact tgaggctttt ggggtgcttg 120
15 ttcagtccat ttgcctgag ggtgcagatt gcccttaacc tcaagggttt ggattatgag 180
gttggtgaag agactttgaa tcccaaaagt gaattgcttc ttaagtccaa ccctgtgcac 240
aagaaaatcc cagttttctt ccatggagat aaagtcatat gtgaatctgc aatcatagtt 300
gagtacatag atgaggtttg gtccaacaat gctctctcca tccttcaca aaatgcata 360
gatcgagcta atgcccgatt ttgggtttct tacatcgatg acaagtggct tacgtccttg 420
20 aaaagtgttc tagcgactga agatgatgag gcaaagaagc tacactttga gcaagcggaa 480
gaagtgcctg agaaggtgga agaagtgttc aacaagtgca gtgaaggga ggcctatttc 540
ggaggagata cgattggatt tgttgacatt ggttttggaa gctttttgag tttcattaga 600
gtctcagaga atatgaatga aagaaaattg cttgatgaaa cgaagtaccc tggtttgacc 660
ctatgggctg aaacttttgc tgcgtatcct gctgtgaagg gccttctgcc agagactgaa 720
25 aagcttgctg agtttgcaaa gattcttcag ctaaaatggg ctgctgcagc tgctgcaaa 780
taaagtgaat caaattaatt gctggatgaa tttcaaaaat tgttggtgca gttatttata 840
tctgaggcta tgtttgttgc aactttatat atttaaaagt caaaataaat gttatgataa 900
tatagtaaaa aaaaaaaaaa 918

```

30 <210> 15

<211> 26

<212> DNA

<213> Artificial Sequence

35 <220>

<223> Description of Artificial Sequence:PRIMER

<400> 15

gcgaagcchc armgagargg hggagg

26

40

<210> 16

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PRIMER

---

5  
    <400> 16  
    cgcaactgaga gaggatcctc gag 23

10  
    <210> 17  
    <211> 27  
    <212> DNA  
    <213> Artificial Sequence

15  
    <220>  
    <223> Description of Artificial Sequence:PRIMER

20  
    <400> 17  
    ttctgggykr astwcityga craiaag 27

25  
    <210> 18  
    <211> 33  
    <212> DNA  
    <213> Artificial Sequence

30  
    <220>  
    <223> Description of Artificial Sequence:PRIMER

35  
    <400> 18  
    gagtcymwkg tsattgttga atacattgat gag 33

40  
    <210> 19  
    <211> 23  
    <212> DNA  
    <213> Artificial Sequence

45  
    <220>  
    <223> Description of Artificial Sequence:PRIMER

50  
    <400> 19  
    cctctcaaac ccatggctca acc 23

55  
    <210> 20  
    <211> 31



&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

---

5 <223> Description of Artificial Sequence:PRIMER

&lt;400&gt; 20

gcgctcgaga gttaggtata cagtatctac c

31

10 &lt;210&gt; 21

&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

15 &lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:PRIMER

&lt;400&gt; 21

cctctcaaac ccatggctca acc

23

20

&lt;210&gt; 22

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

25

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:PRIMER

&lt;400&gt; 22

30 cgcggtacct ccatacaaag aaaatca

27

&lt;210&gt; 23

&lt;211&gt; 24

&lt;212&gt; DNA

35 &lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:PRIMER

40 &lt;400&gt; 23

gagatctgca acaaacatag cctc

24

&lt;210&gt; 24

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

5 &lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:PRIMER

&lt;400&gt; 24

tacaccatgg ctgaaaggga cttg

24

10

&lt;210&gt; 25

&lt;211&gt; 2763

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

15

&lt;220&gt;

<223> Description of Artificial Sequence:SOYBEAN Derived  
nucleic acid sequence P32110

20 &lt;400&gt; 25

ccatacaaac aatacataac taaataactg taattttatg ttttctttca ttgtattgtg 60  
cgttttatat taatgtatta attatgttgt tttttaattt ttattgataa attatttaaa 120  
ctttaaataa aaaacattta aagtaaacta taataaaaaat atatataaca tgtttttaaat 180  
aattaaacat taaaataaaa taataacata tcaataaaaa tcacaaaaaa tataactaatg 240  
25 aaatcaactt aatattattt attatatatt taaatatata ttttttaact tttttatttt 300  
aaaaaaataa aattgttaac attataaaat taaaaaacia atttatgaaa aaaaagggtta 360  
acaaattaaa aacaaactac aaatagtaaa ataacgtttt taaaaaaaaa aggagggttac 420  
gatctcaaaa tcgtatacct catgaaattt taaaaaata agtaagacag ttttacaact 480  
ttaaagtaat aaaacaaaat acgagttttt ttttaatacg actttaaggt cgtaaaaaaa 540  
30 ttatatttac attgaagttg taaataatat ttttacgatt tcagttaaaa aaataagaaa 600  
actaaaaagt cgtaataggt gctaacaact gttacaactt aaccggtttt agataatttt 660  
ttaaaatata ccccaaatga aaaattgctg attattttat cccaatttg taaaaaagca 720  
cctgaatact agaaacaagg gtggcaaaga ttcaaagtaa tgaatttata aaaaaaaaaa 780  
gaaaatattt atataattac acaaaatctt aaaaaatgtt gttattattt actactaatt 840  
35 tataattcga atcgagtata tttaaggcaa agttctctac gcggtttacg taatctctta 900  
catcatttat ttttccacgt ttgtttactc tgaggcatct tcttgatagg ggaaagtttt 960  
ttcatatttt ttctagatcc ttctcatgtg caatgtctcc gcattcatac gcagcaatca 1020  
aaatggaatt aatacacggt agcactttcc catttttttg ttatcgctcc cactattgac 1080  
tatacccatg atatgtatat aggtatatct tttctaattg attacgtatc tgctgaatac 1140  
40 tagctagtcc tgattcctag ctctataaaa ggagaatacc ataggaattc atcacagaca 1200  
aacaacaat ttaccagcta tacttgttcc ttttgaaggt tagaagtgtc acaatacaaa 1260  
caatggcagc tactcaggaa gatgtgaagc ttttggttat tgtgggaagc ccatttgtgt 1320  
gcagggtcca gattgccctt aagttgaagg gagttgaata caaatttttg gaagaaaatt 1380

```

tgggcaacaa gagtgatttg cttctcaa ataccctgt tcacaagaag gttccagtgt 1440
ttgttcacaa tgagcagccc atagcagagt ctttgtgat tgttgaatac attgatgaga 1500
catggaagaa caaccccatc ttaccttctg atccttacca aagagccttg gctcgtttct 1560
gggccaaatt cattgatgat aaggtaactc aacatttcaa aaatcttctt agttttcatg 1620
5 atttgtgctg atttgtcagc aaaacatcac gatgaaatct atatatgtga aatctttctg 1680
gtgtggaata tatatgtgaa atctttgaat atgttagaga actcaaaagt caacagccaa 1740
ccatgatttt tttttaatgt atcaactttt tgcataacaa tattagtgat ttgaaacttt 1800
atggaatcat atactatgat tttgggagaa attttatttt attttactat ttttatctgg 1860
gggtggggga ttctgtcacg tatttgtttc tatataactc gatctaaatt ctgtttgtcc 1920
10 taatcacttt atgaaataat tactaataaa tattgtgatt tgcgaaatca gattgtgggt 1980
gctgtatcga aatctgtttt caggttgat gagaaagagc gtgagaagaa tgttgaagaa 2040
acatatgagg ctcttcagtt tcttgagaat gagctgaagg acaagaagtt ttttgaggga 2100
gaggaatttg ggttggtaga tattgctgct gtcttcatac cattttggat cccaattttt 2160
caggaaatag cagggttgca gttattccac agtgagaaat ttctatact ctacaaatgg 2220
15 agccaagaat tccttaacca cccttttctg caggaagtc ttctctctag agaccactt 2280
tttgctact tcaaagcccg ctatgaaagt cttctgctt caaaatagac ttatttaagg 2340
atatttggtg aacaacttgt gtcttggtga gttattgctg tttgaatttc atgtaaaatg 2400
atactagcta tatgtaaatc ccagaaaaaa aaaaaaaga atcctaggat ctgtttttcg 2460
ttttggccat ttcagtatat aaagaaatta tatttttcga tataaatttt gttgtgaaaa 2520
20 gctttattct tccttcataa aatccttcaa tgtgcataat cttattcgta gagagactta 2580
gagcggctag tagctactac cttgaaattt ttctttta atcgaggaca acgtatatat 2640
tatataataa taattattgc aagttgaaa tcgtgtaagc atgtttcatg actatatgag 2700
ttaacaatat actgtcttgc ctctgcaacc ttcattgatt ctaaaattat tcccttggct 2760
gca 2763
25
<210> 26
<211> 1137
<212> DNA
<213> Artificial Sequence
30
<220>
<223> Description of Artificial Sequence:Mungbean
Sequence U20809
35
<400> 26
gaattaaagt gtttcgatgg gttcaagtca ggaagaagt acccttctgg aattattgga 60
agcccatttg tttgcagggt gaagatagcc ctgaagttga agggagttga atacaaatac 120
gttgaagaaa atttccgcaa caagagtga cagcttctga aatacaaccc agttcacaag 180
aaggttccag tgtttgttca tggtgacaaa ccccttccag agtcccttgt gattgttgag 240
40 tacatcgatg agacatggaa caacaacccc atcttggctt ctgatactta ccagagagcc 300
ttggctcgtt tctgggtcaa attcatcgat gacaagattg tgggtgcttc gtggaaatct 360
gttttcacag ttgatgagaa agagcgtgag aagaatattg cagaaacata tgagagtctg 420
cagtttcttg agaatgagat aaaggagaag aagttctttg gaggagaaga gcttggggtg 480

```

gtagatattg ctgctgtcta tgtagcattt tggatccctt tgattcaaga aatagcagga 540  
ttggagttat tgacaagtga gaaatttcct aatctctaca ggtggagcca agaatttttg 600  
aaccatccaa ttgtcaaaga aagtcctccc cctagagacc cagtttttgc ctttttcaaa 660  
ggacgctatg aaggcctttt ttcttcgaaa tagatttcat gttgtgagag atttagaatt 720

5 tataaggaaa attgtgtgga gtacttagtt aggatttggg ttcaaaatta tggttgaagt 780  
tgaatcctag gatttgcgca tgtcaaacaa ataacctggg attgttcgtg ttgatatttt 840  
actatttcaa tcaataaatt atgcagcttc tiaccgagtt aacattcgat cgaaataagg 900  
accaacaaga ttaagtaagg ctgcattatt tgtctttttg ttaaattaga tattagtatg 960  
caccaaaaag tgagtatttc cttacagaag ctttttaaat attaagtagt taattccata 1020  
10 ggtctacat tatagctcaa gttatataca tattatgggt gccattctct actcaacaat 1080  
tatgactata aaatcttgtg gttataatgc cacgaacaag tgaactatct cacttca 1137

<210> 27

<211> 2038

15 <212> DNA

<213> Artificial Sequence

<220>

20 <223> Description of Artificial Sequence:Tobacco  
sequence Q03663

<400> 27

ctcgaggatt tcaaactcta gtttactaa aacttgagct ttcttttcca ctaatgtcga 60  
aaaacgaaat aaacataagc tatttacaaa aaataaaaaa atactccatt tgaatctaaa 120  
25 gtcaagtcgt gattgggata agaaaataga aattttattta tactccagat caagccgtga 180  
ttggaatgag ataatagaaa agtatgatag tacatgagta acatcaagtt ggaaattaag 240  
ggaaggaaat tagagaaaga actgaagaat atccaaatat tctttacgtc caaatttgat 300  
agttatttaa cgtcacgcag atgacggcca tgttcaagtt ttccacaaat attgagaaaa 360  
gaaagaagaa gacacaaact gtgttttgta ttattatagt tttttctttt agagaattga 420  
30 ttgtacatat aagaaatata atataagatt tagaaataag attattagaa aaatcaaaca 480  
tcaaagtatt tatttttaaat tctttttcca atggacattc ccattctgaa aaaaaagaga 540  
tataaatatg gaagtaaaaa ttaatcagat cgttaaatgt agaaaatatt aattaacaca 600  
ttaaccataa ccagtctact ttatttaaca aaaagcacat ctgaratarc aaaaaagtgt 660  
ttaacttcat gcattgacaa tttaaaatta ttttgcaaca tcgggtaaaa ctattttaca 720  
35 acaattggta actgcatata taagtttaat atggtaacct agaaaatagg ataaattatc 780  
tataacagga tatattacat tgatattacc atgtcaaaaa atttagtaag tacatgaata 840  
atcaccgtga aatcttcaag atttctccta taaataccct tggtagtaaa tctagttttt 900  
ccattcaaga tacaacattt ctctatagat catgggattt gttctctttt cacaattgcc 960  
ttcattttctt cttgtctcta cacttctctt attcctagta atatcccact cttgccgtgc 1020  
40 ccaaaattct caacaagact atttgatgc ccataacaca gtcgtgcag atgtaggtgt 1080  
agaacctttg acctgggacg accaggtagc agcctatgcg caaaattatg cttcccaatt 1140  
ggctgcagat tgtaacctcg tacattctca tggtaatac ggcgaaaacc tagctgaggg 1200  
aagtggcgat ttcattgacg ctgctaaggc tgttgagatg tgggtcgatg agaaacagta 1260

```

ttatgaccat gactcaaata cttgtgcaca aggacaggtg tgtggacact atactcaggt 1320
ggtttggcgt aactcgggtc gtgttggtatg tgctaggggt cagtgtaca atggagaata 1380
tgttgtctct tgcaactatg atcctccagg taattataga ggcgaaagtc cataactaatt 1440
gaaacgacct acgtccattt cacgttaata tgtatggatt gttctgcttg atatcaagaa 1500
5 etttaaataat tgetctaaaa ageaaettta agtcaagtat atagtaatat tactatattt 1560
gtaatcctct gaagtggatc tataaaaaaga ccaagtggtc ataattaagg ggaaaaatat 1620
gagttgatga tcagcttgat gtatgatctg atattattat gaacactttt gtactcatac 1680
gaatcatgtg ttgatggctt agctacttgc gatattacga gcaaaattct taactacatg 1740
ccttaggaac aagcttacac agttcatata atctactaga gggccaaaaa catgaaaatt 1800
10 accaathtag atggtaggag gatattgaaa gtggagcagc tagttttaat aactgaccgt 1360
tagtcttaaa attgacggta taaaaatatt tacataatca ggtcatttat aaggtaatta 1920
taggtaaata tttatgacga attctcaata gtaatctgaa aaaaaattgt aactaaccta 1980
ttatactaaa actactataa taggttagat tacattaatc atgtcattag aagatctt 2038

```

15 <210> 28

<211> 2796

<212> DNA

<213> Artificial Sequence

20 <220>

<223> Description of Artificial Sequence:Potato- Derived  
nucleic acid sequence P32111

<400> 28

```

25 aagcttacat tcaacgtgtt gctgcttcaa aataagggtc tttacaaaat ttaaaaaata 60
taaagaagaa ttttgaattt tataattaaa gccgtcagaa aggacttgac ctttgaagcc 120
aaccacaactc aatattagaa aatcaaaaat tttagtgcac tcattttataa aaaaaaaaaa 180
aattacttat gcagttcttg aaccctttgt gagacgagag ggagttgctc ggatggtaag 240
cacccttcac tttcaaccgg aaggttgcca gtttgagtca ccaacggagc aaaaagggtg 300
30 ggagctccta gaaagggtaa aaaaaaaaaa aaaaattaat aaaaaaatat cctttatgaa 360
atctctcatt ccgctactgc acttctcccc tgatcttctt cgtgttttca attattaatt 420
ctatattcat gacaccatgt gatgtttctc tgggtagtcc taaaaataga ggtattgaaa 480
attatgttgt ttctctctgg tctatttact tttcttgtgt actttattgt atttcatatt 540
gttaattttt ggcttcggtt tataacatgt tattagcaca aaactttaat catatcgagt 600
35 taaactttta attttgctta tcaacgtaa agacaagata tgtgatcggc atgtataact 660
atgttttaat taggtataat acataaatat ttcttttaat tttatctcat tttatattta 720
tgtcgtttta ctttgaggcg tacaagtgtc tacttgatgt ccagtagggt atcaaagctg 780
gcatgcttag ttttactttc caaattgaaa tttatattag aattgaattc aggaagaatt 840
ttgtaggttc aactaaatta tatatatata tataaaaaaa taaaattat tagacgcttc 900
40 gactatttac ttacttttaa atttgaattt tcgtacgaat aaaattattt gtcagagaaa 960
agtcttttag ctattcacat gctaggaagt ttacttttg gtggatcagt gattgtatat 1020
tatttaatat atatcaattt tctcatcaa ctgaaaatga aagataaaat taatattaaa 1080
aactccattc attttaattt attgtcatgt tttgacttga tccaaaatct aacaatttaa 1140

```

```

aagggttttaa atttttgtgc ttttttttaa attaaaaata tgtcaaatat attaaaaatat 1200
atttttttaa ttttatacta aaaaacatgt cacatgaata tttgaaatta taaaattatc 1260
aaaaataaaa aaagaatatt tctttaacaa attaaaaatg aaaatatgat aaataaatta 1320
aactattcta tcattgattt ttctagccac cagatttgac caaacagtgg gtgacatgag 1380
5 cacataagtc atetttattg ttttttatta cteactccaa aaatataggg aatatgttta 1440
ctacttaatt tagtcaaata taattttata ttagaataat tgaatagtca aacaagaaac 1500
tttaatgcat ccttattttt tctctataa aaaaaagact agacaccaag ggagaccaac 1560
cacacataat taagatggca gaagtgaagt tgcttggtct aaggtatagt ccttttagcc 1620
atagagttga atgggctcta aaaattaagg gagtgaata tgaatttata gaggaagatt 1680
10 tacaaaataa gagcccttta cttcttcaat ctaatccaat tcacaagaa attccagtgt 1740
taattcacaa tggcaagtgc atttgtgagt ctatggcat tcttgaatac attgatgagg 1800
catttgaagg cccttccatt ttgcctaaag acccttatga tcgcgcttta gcacgatttt 1860
gggctaaata cgtcgaagat aaggtatatt gcttttaagt tattccaatt gattgaaaag 1920
tttgttttag ttacgttatt acatatactt taggtctcat gctttttaat aatcttttat 1980
15 aaaattcgac taagacgaac ttctcgtata gtcaacaata ctaacatatt tgtctagtag 2040
ttggtttaga aataagttat ccgaatatta aattctggat aagtaatgaa taccatattt 2100
gatagttgat ttggagataa attattcgtg tataaaatta atatgatatt tgatttgcaa 2160
tttagaaata cataactatt ttatatgcat agatccatta taactaattg atatattatt 2220
aatatctgta taactctaac cagctatcga aacgagtcaa cgaaccttat taagttttgt 2280
20 ttgttgggca gggggcagca gtgtggaaaa gtttcttttc gaaaggagag gaacaagaga 2340
aagctaaaga ggaagcttat gagatgttga aaattcttga taatgagttc aaggacaaga 2400
agtgccttgt tggtgacaaa tttggatttg ctgatattgt tgcaaatggt gcagcacttt 2460
atlttgggaat tcttgaagaa gtatctggaa ttgttttggc aacaagtga aaatttccaa 2520
atltttgtgc ttggagagat gaatattgca cacaaaacga ggaatatatt ccttcaagag 2580
25 atgaattgct tatccgttac cgagcctaca ttcagcctgt tgatgcttca aaatgagtat 2640
acctcaagtg aatttcaaga ttttgtgtgg caataaaaaat tgagtttttg taaattcaat 2700
tgaaatatat taaagttgca tgttataaga tttatcttta tttcactagt taatataaat 2760
tttgattca cgtataaata aaagtattgt taagag 2796

```

30 <210> 29

<211> 1289

<212> DNA

<213> Artificial Sequence

35 <220>

<223> Description of Artificial Sequence:Arabidopsis

P46421

<400> 29

```

40 aagtcaagg acacatgctc cagggataag gcaagggttag gaattaggac acatctccac 60
ttactaaaaa agagataaaa aaaaattgta tagggaacgt tataaatatg ttgtaaagtc 120
aacatctgtt tcttctaga ctcttcgcat ttacatcaca ctgccgacca tataaaacgg 180
caaagttcgt cgtcgtttta tcacaagacc atcaacacca taaggctata aatccaagct 240

```

```

aaaaggtagt gattaactcc acaaaaccag aaaaactaca tttctaacat atagaagaaa 300
cagagaaaaa gagagagaga cccctaattg ctgagaaaga agaagtgaag cttttgggga 360
tatgggcgag cccttttagc cgtcgggtcg agatggctct caaactcaaa ggcataccgt 420
acgagtacgt ggaagagata ctggagaaca aaagcccttt gcttcttgct cttaaccta 480
5 ttcacaagaa agtccctggt cttgtccaca atggtaaaac cattctcgag tctcatgtga 540
ttcttgaata catcgatgag acttggccac aaaatccaat tctccctcaa gatccttatg 600
aaagatccaa agctcgtttc tttgctaaac tcgtcgatga acaggtaatt gaattgggtc 660
aaaattgcat gtcaaataat aaacaattgg ttctgctttg ttaatttatc aaacaagtaa 720
ttttctatta acattagcga ttatatgtct ctgtcattgt agattatgaa cgtgggggtt 780
10 atatcaatgg caagagcaga cgagaaagga agagaagttt tagccgagca ggtaagagaa 840
ctgattatgt atctcgagaa agaacttgtc ggaaaagatt acttcggagg caagactgtc 900
ggattcttgg actttgtcgc cggaagtta attccgtttt gtttgagag aggttgggaa 960
ggaataggat tggaagtgat tacagaggag aagtttccag agttcaagag atgggttagg 1020
aatgttgaga aggttgagat tgttaaagat tgtgttccac caagagagga acatgtagaa 1080
15 cacatgaact atatggcaga gagagtgaga tcttcttaag aaaacaaatc atgtttagtt 1140
cttgatcatg caatgtttgt atggttatgt tgttgtttat tttattgaat atctttgtat 1200
gttggtgtgt tgagaagtga ggttttatca tcatctctca cgttatctta tttggtcca 1260
gccactattt agaattaatg gtaaagctt 1289

20 <210> 30
    <211> 1339
    <212> DNA
    <213> Artificial Sequence

25 <220>
    <223> Description of Artificial Sequence:Arabidopsis
        Genomic sequence

    <400> 30
30 gaattaattc ttgacgaagc atgggggtgt cttcagaaaa tgaatggatc aagtcaaggt 60
   acacatgctc cagggataag gcaaggttag gaattaggac acatctccac ttactaaaaa 120
   agagataaaa aaaaattgta tagggaacgt tataaatatg ttgtaaagtc aacatctgtt 180
   tccttctaga ctcttcgcat ttacatcaca ctgccgacca tataaaacgg caaagttcgt 240
   cgtcgtttta tcacaagacc atcaacacca taaggctata aatccaagct aaaaggtagt 300
35 gattaactcc acaaaaccag aaaaactaca tttctaacat atagaagaaa cagagaaaaa 360
   gagagagaga cccctaattg ctgagaaaga agaagtgaag cttttgggga tatgggcgag 420
   cccttttagc cgtcgggtcg agatggctct caaactcaaa ggcataccgt acgagtacgt 480
   ggaagagata ctggagaaca aaagcccttt gcttcttgct cttaaccta ttcacaagaa 540
   agtccctggt cttgtccaca atggtaaaac cattctcgag tctcatgtga ttcttgaata 600
40 catcgatgag acttggccac aaaatccaat tctccctcaa gatccttatg aaagatccaa 660
   agctcgtttc tttgctaaac tcgtcgatga acaggtaatt gaattgggtc aaaattgcat 720
   gtcaaataat aaacaattgg ttctgctttg ttaatttatc aaacaagtaa ttttctatta 780
   acattagcga ttatatgtct ctgtcattgt agattatgaa cgtgggggtt atatcaatgg 840

```

caagagcaga cgagaaagga agagaagttt tagccgagca ggtaagagaa ctgattatgt 900  
 atctcgagaa agaacttgtc ggaaaagatt acttcggagg caagactgtc ggattccttg 960  
 actttgtcgc cggaagtta attccgtttt gtttgagag aggttgggaa ggaataggat 1020  
 tggaagtgat tacagaggag aagtttccag agttcaagag atgggttagg aatttgaga 1080

5 aggttgagat tgttaaagat tgtgttccac caagagagga acatgtagaa cacatgaact 1140  
 atatggcaga gagagtgaga tcttcttaag aaaacaaatc atgttttagtt cttgatcatg 1200  
 caatgtttgt atggttatgt tgttgtttat tttattgaat atctttgtat gttgtgtggg 1260  
 tgagaagtga ggttttatca tcatctctca cgttatctta tttgggtccca gccactatct 1320  
 agaattaatg gtaaagctt 1339

10

&lt;210&gt; 31

&lt;211&gt; 968

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

15

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Papaya AJ000923

&lt;400&gt; 31

20 tagaactagt ggatcccccg ggctgcagga attcggcacg agagatttta tcttttagga 60  
 gctccgtttt acaacaatgg cggacgaggt tgttctcttg gatttctggc caagcccttt 120  
 tggaatgaga atcagaatcg ctttagccga gaaggggtatt cactacgagt acaaggaaga 180  
 gaatctgaga aacaagagtc ccttactcct gcagatgaac ccggtacaca agaaaatccc 240  
 ggttctcatc cacaatggta aacctatctg tgagtctttg atccagattc agtacataga 300  
 25 tgaggtatgg agcgacaagg ctctctgct tccctctgat ccttatcaga gagctcaagc 360  
 caggttctgg gctgactatg ttgacaagaa gatgtatgaa gctgggagga gagtttggac 420  
 gactaaaggg gaagaacagg agggggccaa gaaagagttc atagaaatct tgaagacttt 480  
 ggaggggagaa cttgggggaga agccttattt tgggtggggaa agttttgggt atgtggattt 540  
 gacttttatc ccattctaca cttggttcag tgtgtatgaa agttttggga agatgagcat 600  
 30 agaggcagaa tgccccaagt tgttttagttg ggtgaaaagg tgtttggaga aggagagtgt 660  
 ttcaaaatct ctgcctgatc aagataaggt atacggcttc gttttggaac tcaggaaggc 720  
 tcttgggatt tgagtttttt gagagacttc aaaatccttg ttccatttcc attaggggtt 780  
 gtctccaag tattgattaa aaaaggttct ggatcaacta ctttatttgt ctagtctttt 840  
 actgttgtat tggaataaag ggtgtcctt ttgtggatta ggtgagattt ctatcaatat 900  
 35 tgtggtgacg tcaatttctt gtgtgttgta ggcaaatcat atttgaataa aatctttctt 960  
 tcatatgt 968

&lt;210&gt; 32

&lt;211&gt; 1040

40 &lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;



<223> Description of Artificial Sequence: Spruce AF051214

<400> 32

```

gaaaagacag ataatagaa agttagagac agttacaata cattttcaga gcaatcgca 60
5  tggaggcctg tggagaagag gcgcaagtga agctgttggg tgggaatata agtccctttg 120
   tgctgcggtt ccgtatagca cttgctctta aagggatcga ttacgagttc atcgaagaga 180
   acatgcaaaa taagagccat ctgctgctgc aatcgaaccc tgtaacaag aagattccag 240
   tgctcatcca caatggaaag cctgtttgtg agtccatgat tattgtgcag tacatagacg 300
   aggcattgga caggaaggcg ccagttctaa tgcccaaaga tccttatgac cgagccattg 360
10  cccgcttctg ggctgccttt gtagacgata agctgctgcc atgccttcgg ggagttttca 420
   agggccaggg agagcagcaa cagaaagcgt tggaagaatc gggggcaagc tttcttttac 480
   tggaggaggc tctgcgaacg agccactgct tctcgggaaa accgtatttc ggaggagatg 540
   agatcggctt tcttgacatc gcattgggtg gtatgttagc atttgtcaaa gccctcgaga 600
   aggttactaa tttagtttta atagaccagg agaagatgcc gctgttaagc acatggatga 660
15  atcgattctg tgaggccgat ggagtgaag acgttatgcc ggatccggcc aagttgcagg 720
   aatttatatc cgccatcaga gtcagattta catcaccacc tgctgccaat taggggaagc 780
   cattcgcca ttaaatggat gttatcgtcc gcattgtttt tggttttatg ctgtcagttt 840
   gaatgttgtt atgctttttg aattgttgtt gttaaatggg aataattcta tcgccccaatc 900
   tagcaccgtg tgattcgcta tcagttctca ccgtgttcca tgtaacttcg atttcatgat 960
20  tttgggagat agaacaaaaa ttcattgaaa tgtgtgttag tgttttatat ttgaaaaggg 1020
   ttgatttgc agagaatgga                                     1040

```

<210> 33

<211> 902

25 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Wheat AF004358.

30

<400> 33

```

ggcacgaggc aagcacaagc agctaaagca atctcagtgg tcttcaaaac acacacacac 60
   aactgacac acatcgatcg aggtagttag agatggccgg aggagatgac ctgaagctgc 120
   tcggcgcatg gccaagccca tttgttacca ggggtgaagct ggcgctcgcc ctgaagggcc 180
35  tgagctacga ggacgtggag gaggacctgt acaagaagag tgagcttctc ctcaagtcca 240
   acccggtgca caagaagata cccgtgctca tccacaacgg agccccggtc tgcgagtcca 300
   tgatcattct ccagtacatc gacgaagtgt tcgccagcac cggcccgtcc cttcttccag 360
   cggaccctta cgagcgcgcc attgctcgct tctgggtggc ttacgttgac gacaagctgg 420
   tagcccatg gaggcagtgg ttgaggggca agacagagga ggagaaatcc gagggaaaga 480
40  agcaggcggt cgccgcggtg ggggtcctcg aaggggccct gagggagtgc tccaaggag 540
   ggggcttctt cgggtggcgc ggcgtcgggc tcgtcgacgt tgcgctggga ggcgtgctgt 600
   cgtggatgaa ggtgaccgag gcgctgtctg gtgacaagat tttcgacgcc gccaagactc 660
   cgctcctggc cgcattgggtg gagcgcttca ttgagctcga cgcggccaag gccgccttgc 720

```

```

cggacgtggg caggctgctt gagtttgcca aggcacgaga ggctgccgct gcagcgtcca 780
agtgagccgc cagcacatat ccagaataat taaaatttgt tattttaaat gttgttcgtt 840
cggctgcttg atgtaataat gtagtaactg atgtcgtcca tttaaaaaaa aaaaaaaaaa 900
aa                                                    902

```

5

```

<210> 34
<211> 1127
<212> DNA
<213> Artificial Sequence

```

10

```

<220>
<223> Description of Artificial Sequence: Spruce AF051238

```

```

<400> 34

```

```

15  gacacatata actcacaggc aaaaaaatat tcaattacaa tacattttcc gtgcaatggc 60
    gacggaggct tgtggagaaa aggggcaagt gcagctgtta ggtgggagtc tgagttcctt 120
    cgtgctgcgg gttcgcatag cacttgctct taaaggcatc gattacgagt icatcgaaga 180
    gaacatgcaa aataggagcc agctgctgct gcaatcaaac cctgttcaca ggaagattcc 240
    agtgcttata cacaatggaa agcccgtttg tgaatccatg attattgtgc agtacatcga 300
20  tgaggcatgg gacactaagg cacccaatct tatgcccaaa gatccatatg acctagccat 360
    tgcccgttcc tgggctgcct ttgtagacga taagctcggt ccatgtatgc ggagagtttt 420
    cgctggccat ggcgagcagc tacagaaaga agcggaagat ctggttacaa actttcattt 480
    gatagaggaa gctctgcgaa ccaacagctg cttctcagga aaagcgtatt ttggagggga 540
    taagataggc ttgcttgaca tcgcattggg tggtagtttg gcggttctca aaggccctga 600
25  gaaggctacc gataccgtta taatagatcc ggagaagatg ccgttgctga gcgcatggat 660
    ggaccgattt tgtcaatcca atggagtga agaagtaatg cccgatccgg ccaagcagct 720
    ggaatctcta tcagctagga gagccagact tgcatcacct gctggcaatt agggcaagcc 780
    atgtcggcct tataaactga ggatagacag atggattata aacttattat tcgtagtact 840
    tgtcctttta ttcattgtgt cagcttcagc gttttaattc ttgctgtttt atgtgaataa 900
30  gtctgaataa tgtttgggtg aatctgcct gtactatagc tggcattcac ctgtttattg 960
    tacgctgatt tagtttgaac aagttttggt gaatctcccc tgtactgaag ctggcattcc 1020
    cctgttcaat gtgcgctgat ttagtttgaa taagtttttg atgaatctcg cttgtactgt 1080
    agctatgtgc gatgatTTTT aatgccatag aaacgagaat gaaatgc 1127

```

35

CLAIMS

1. A protein comprising the amino acid sequence depicted as SEQ ID No. 1 or a protein  
variant having at least about 70% identity therewith wherein the said protein or  
5 variant is capable of catalysing the addition of Beta-alanine onto gamma  
glutamylcysteine.
2. A protein variant according to claim 1 having a  $K_m$  for Beta-alanine which is less  
than the said variants  $K_m$  for glycine when calculated using the same method.
- 10 3. A protein variant according to claim 2 having a  $K_m$  for Beta-alanine which is less  
than or equal to about 0.8mM and a  $K_m$  for glycine which is higher than 0.8mM  
when calculated using the same method.
- 15 4. A protein variant according to claim 2 or 3 which variant comprises an amino acid  
sequence selected from the group depicted as SEQ ID No. 2, 3, 4 or 5.
5. A polynucleotide comprising a region encoding the protein or protein variant  
according to any one of claims 1 to 4.
- 20 6. A polynucleotide according to claim 5 which comprises the sequence depicted as  
SEQ ID No. 6.
7. A Glutathione-S-transferase comprising the amino acid sequence depicted as SEQ ID  
25 No. 7, 8, 9 or 10.
8. A polynucleotide comprising a region which encodes at least one of the amino acid  
sequences according to claim 7.

9. A polynucleotide sequence which is the complement of one which will hybridise to the polynucleotide according to claim 8 under stringent conditions and which polynucleotide sequence still encodes a Glutathione S transferase.
- 

- 5 10. A polynucleotide according to claim 9 which comprises the sequence depicted as SEQ ID No. 11 or 12 or 13 or 14.
11. A polynucleotide sequence which is the complement of one which binds to SEQ ID No. 11 or 12 or 13 or 14 at a temperature of between 60°C and 65°C in 0.3 strength  
10 citrate buffered saline containing 0.1% SDS followed by rinsing at the same temperature with 0.3 strength citrate buffered saline containing 0.1% SDS wherein said polynucleotide sequence still encodes a functional Glutathione S transferase enzyme with the *proviso* that the said polynucleotide sequence or the amino acid  
15 sequence which it encodes, is not a sequence selected from the group of sequences listed under accession number; P32110, U20809, Q03663, P32111, P46421, AJ000923, AF004358, AC000348 or AF051238.
12. A DNA construct comprising in sequence a plant operable promoter operably linked to a polynucleotide according to either of claims 5 or 6 operably linked to a  
20 terminator region.
13. A DNA construct comprising in sequence a plant operable promoter operably linked to a polynucleotide or polynucleotide sequence according to any one of claims 8 to 11, operably linked to a terminator region.  
25
14. A DNA construct comprising a first region comprising in sequence a plant operable promoter operably linked to a polynucleotide according to claim 5 or claim 6 and a terminator region and a second region comprising in sequence a plant operable promoter operably linked to a polynucleotide or polynucleotide sequence according  
30 to any one of claims 8 to 11 and a terminator region.

15. A DNA construct according to claim 14 wherein the said first region comprises a polynucleotide encoding the amino acid sequence depicted as SEQ ID No. 1 and the second region comprises a polynucleotide encoding the amino acid sequence depicted as SEQ ID No. 10.
- 
- 5 16. A DNA construct according to any one of claims 12 to 15 which further comprises a third region which provides for the production of a protein which acts as a selectable marker.
- 10 17. A DNA construct according to any one of claims 12 to 16 wherein the plant operable promoter is selected from the group consisting of: CaMV35S, FMV35S, NOS, OCS, Patatin, E9, alcA/alcR switch, GST switch, RMS switch, oleosin, ribulose bisphosphate carboxylase-oxygenase small sub-unit promoters.
- 15 18. A method of providing plants which are resistant and/or tolerant to an agrochemical comprising:  
(a) inserting into the genome of plant a polynucleotide or a polynucleotide sequence according to any one of claims 8 to 11 or a DNA construct according to any one of claims 13 to 17; and  
20 (b) regenerating morphologically normal fertile plants or plant parts therefrom; and  
(c) applying to said plants or plant parts an amount of said agrochemical which is phytotoxic to control like plants and selecting those plants or plant parts which are resistant to said agrochemical.
- 25 19. A method of providing plants which are resistant and/or tolerant to an agrochemical comprising:  
(a) inserting into the genome of a plant which plant provides for the production of a functional Glutathione S transferase, a polynucleotide according to either of claims 5 or 6 or a DNA construct according to claim 12; and  
30 (b) regenerating morphologically normal fertile plants or plant parts therefrom; and

(c) applying to said plants or plant parts an amount of said agrochemical which is phytotoxic to control like plants and selecting those plants or plant parts which are resistant to said agrochemical.

---


- 5     20.     A method according to claim 18 or 19 wherein the said agrochemical comprises acifluorfen and/or chlorimuron-ethyl and/or fomesafen and/or acetochlor and/or metolachlor.
- 10     21.     Plants or plant parts obtained according to the method of any one of claims 18 to 20.
- 10     22.     Plants or plant parts according to claim 21 wherein said plants or plant parts are selected from the group consisting of: melons, mangoes, soybean, cotton, tobacco, sugarbeet, oilseed rape, canola, flax, sunflower, potato, tomato, alfalfa, lettuce, maize, wheat, sorghum, rye, bananas, barley, oat, turf grass, forage grass, sugar cane, pea,
- 15     field bean, rice, pine, poplar, apple, peaches, grape, strawberries, carrot, lettuce, cabbage, onion, citrus or nut plants.
- 20     23.     Use of a polynucleotide according to any one of claims 5,6, 8 or 10 or a polynucleotide sequence according to claim 9 or 11 or a DNA construct according to any one of claims 12 to 17 in a method of producing plants which are resistant and/or tolerant to a herbicide comprising acifluorfen and/or chlorimuron-ethyl and/or fomesafen and/or acetochlor and/or metolachlor.
- 25     24.     Herbicide resistant plants obtained by the method of any one of claims 18 to 20.
- 25     25.     A method of selectively controlling weeds in a field said field comprising crop plants and weeds said method comprising applying to said field an agriculturally acceptable formulation of an agrochemical comprising acifluorfen and/or chlorimuron-ethyl and/or fomesafen and/or acetochlor and/or metolachlor or a functionally related
- 30     analogue thereof, characterised in that the said crop plants are the plants according to claim 21, 22 or 24.

26. Use of an agrochemical comprising acifluorfen and/or chlorimuron-ethyl and/or fomesafen and/or acetochlor and/or metolachlor or a functional analogue thereof to selectively control weeds in a field which field comprises crop plants and weeds comprising applying to said field an agriculturally acceptable formulation of said agrochemical in an amount which is sufficient to be phytotoxic to said weeds but not said crop plants characterised in that said crop plants are the plants according to claim 21, 22 or 24.
- 
27. A protein comprising the sequence depicted as SEQ ID No. 7 or a protein variant having a Smith-Waterman score greater than 1094 in the SWISSPROT database calculated using the FASTA3 algorithm wherein the said protein variant still encodes a Glutathione-S-transferase.
28. A protein comprising the sequence depicted as SEQ ID No. 8 or a protein variant having a Smith-Waterman score greater than 619 in the SWISSPROT database calculated using the FASTA3 algorithm wherein the said protein variant still encodes a Glutathione-S-transferase.
29. A protein comprising the sequence depicted as SEQ ID No. 9 or a protein variant having a Smith-Waterman score greater than 671 in the SWISSPROT database calculated using the FASTA3 algorithm wherein the said protein variant still encodes a Glutathione-S-transferase.
30. A protein comprising the sequence depicted as SEQ ID No. 10 or a protein variant having a Smith-Waterman score greater than 766 in the SWISSPROT database calculated using the FASTA3 algorithm wherein the said protein variant still encodes a Glutathione-S-transferase.





FIGURE 1





2/3

2/3

FIGURE 2







FIGURE 3



18/10/00

Perfektion der  
Zentral-Abrechnung

---